



*Streamers: A transition from Corona to Jet.*

*VLBI Discovery of Coronal Streamers in  
a Stellar System*


Maria Massi

**Max Planck Institut  
für Radioastronomie  
(Bonn, Germany)**

## Solar corona



Jardine et al. 2006



von Rekowski and Brandenburg 2004

# Coronae


are a general phenomenon present among a wide variety of objects.

- **Sun / cool stars**

- **Young stellar objects** generally show highly elevated levels of coronal activity.


- **Accretion disks**, which exist around protostars, degenerate stellar objects, and supermassive black holes, can be surrounded by a corona.



The solar corona has been observed / studied in greatest detail. How different are the other coronae ?





In the solar dynamo, i.e. the physical process that generates the Sun's magnetic field, a toroidal field is created at the tachocline.....

In a second stage, tubes of toroidal flux emerge to the surface (because of magnetic buoyancy) creating the **coronal loops** (i.e. the building blocks of the corona)





the tachocline is missing in fully convective objects



In order to investigate


the structure of the corona in  
fully / nearly fully convective objects

we performed a series of observations on  
weak-line T Tauri stars

# Weak-line T Tauri Stars

are pre-main  
sequence objects,



where the radiative  
zone either is still  
missing or is deeply  
embedded and  
therefore  
corresponds to a  
very different  
situation with  
respect to the thin  
convective shell of  
the Sun.



# V773 Tau A

a binary system

Effelsberg 100-m telescope and VLA






Massi, Menten, Neidhöfer 2002

Periastron passage: Large flares

Clustering of energetic flares around the periastron passage


# X-rays

SUN:





[/www.aoc.nrao.edu/intro/solar.flare.gif](http://www.aoc.nrao.edu/intro/solar.flare.gif)

## Radio wavelengths



In V773 Tau A the relationship between intensity of the flare occurrence and distance of the two stars indicates another new mechanism of magnetic interaction: The interaction of coronae.



Magnetic reconnection is taking place far out from the stellar surfaces, where the two coronae interact one with each other.

$$15 R < H < 26 R$$

# X-ray observations

## V773 Tau A

Skinner et al. (1997)  
Tsuboi et al. (1998)


Size of  $\sim 1$  stellar radius



Feigelson et al. (1994)  
radio variability

# Radio Observations


Size of  $> 15$  stellar radii



observed a steady X-ray flux combined with


**Radio and X-ray emission come from spatially separated regions**

# The Solar Corona Observed in X-rays




Coronal loop: Closed magnetic lines

Coronal holes: **Open magnetic field lines**




Open field lines converge above the top of the closed loops creating the “helmet”




helmet  
streamer

Streamer


cusps  
(helmet)



... and expand outward from the Sun (streamers) up to 30 solar radii.



Massi, Forbrich, Torricelli, Menten,  
Neidhöfer, Misuri, Bertoldi 2006



The flare decay (e-folding time of  $2.31 \pm 0.19$  hours) corresponds to a slow leakage of relativistic electrons trapped between two mirror points:

one close to the star (at  $2-5 R_{*}$ ) i.e. the helmet  
and the other  
at the top of the streamer (at  $10-20 R_{*}$ ) i.e. where the two streamers interact  
with each other.

# Effelsberg 100m-telescope




VLBA

Cambridge/MERLIN UK



Effelsberg (DE)



Onsala Space Obs (SE)



Bank (UK)



WSRT (NL)



Torun (PL)



DSN Robledo (ES)



Noto (IT)



Medicina (IT)



# VLBA+Effbg

● 📷 📄 🕒 cm

**Beam size 1.3 x 0.5 mas**


Apoastron:  
52 R=3.6 mas

Periastron:  
30 R=2 mas

**1 mas < H < 1.8 mas**

**7days (A-G)**

**5 hours each**



Boden et al. (2007), Massi et al. (2008)

11 March 2004 (A)

V773 Tau A  
VLBA+EB  $\lambda$ 3.6cm

12 March 2004 (B)

a b c d


13 March 2004 (C)

14 March 2004 (D)

15 March 2004 (E)

16 March 2004 (F)

17 March 2004 (G)



Two distinct structures appear in the radio images .

They happen to be associated to the secondary and primary star of the system.

Massi, Ros, Menten, Kaufman-Bernado, Torricelli-Ciamponi, Neidhöfer, Boden, Boboltz, Sargent, Torres (2008)

2 mas

11 March 2004 (A)

V773 Tau A  
VLBA+EB  $\lambda$ 3.6cm

12 March 2004 (B)

a b c d


13 March 2004 (C)

14 March 2004 (D)

15 March 2004 (E)

16 March 2004 (F)

17 March 2004 (G)




Two distinct structures appear in the radio images .

They happen to be associated to the secondary and primary star of the system.

Massi, Ros, Menten, Kaufman-Bernado, Torricelli-Ciamponi, Neidhöfer, Boden, Boboltz, Sargent, Torres (2008)



2 mas

Plot file version 1 created 25-FEB-2007 23:03:48  
CONT: V773TAUA IPOL 8421.490 MHZ V773TAUA B.ICL001.1



Center at RA 04 14 12.9198350 DEC 28 12 12.199470  
Cont peak flux = 4.7014E-04 JY/BEAM  
Levs = 1.000E-04 \* (-1, 1, 1.500, 2, 2.500, 3, 4, 5)

15 R < H < 26 R



- Electrons, relativistically accelerated in the collision, emit synchrotron radiation, making the streamers „visible“ in the radio band
- Simulations predict the brightness peak of optically thin emission to be, where the magnetic field is stronger (i.e. at the mirror points) (references in Zhou et al. 2005)


11 March 2004 (A)

V773 Tau A  
VLBA+EB  $\lambda 3.6\text{cm}$

12 March 2004 (B)

a b c d

13 March 2004



12 March 2004 (B)

The two features, extended  $18R$  each and showing both mirror points are almost lined-up indicating two helmet streamers shortly after a collision.

11 March 2004 (A)

V773 Tau A  
VLBA+EB  $\lambda 3.6\text{cm}$


12 March 2004 (B)

a b c d

13 March 2004 (C)

14 March 2004 (D)

15 March 2004 (E)



12 March 2004 (B)

One image (run -E) taken a few hours after a flare (second the total flux monitoring at the 100-m Effelsberg telescope) show one elongated fading structure substantially rotated with respect to those seen in the B run, as expected for an helmet anchored on a rotating star

11 March 2004 (A)

V773 Tau A  
VLBA+EB  $\lambda 3.6\text{cm}$


12 March 2004 (B)

a b c d

13 March 2004 (C)

14 March 2004 (D)

15 March 2004 (E)



One image (run -E) taken a few hours after a flare show

One elongated fading structure substantially rotated with respect to those seen in the B run, as expected for an helmet anchored on a rotating star ( $P \sim 2$  days)

11 March 2004 (A)

V773 Tau A  
VLBA+EB  $\lambda 3.6\text{cm}$

12 March 2004 (B)

a b c d

13 March 2004 (C)


14 March 2004 (D)

15 March 2004 (E)

16 March 2004 (F)

17 March 2004 (G)

2 mas




The same rotation is seen in run G for the streamer associated to the other star.

# Conclusions:

The factor triggering periodic flares in V773Tau is magnetic reconnection between streamers belonging to the two different stars.

Interbinary collisions in a weak-line binary T Tauri system ----->>




1.

**Helmet streamers are present also on fully/nearly fully convective objects**

# Phenomena associated to helmet streamers ( Schwenn 2006 )

- Slow wind
- "Blobs" or "Plasmoids"
- Violent coronal mass ejection



# *Bridging the Gap: Corona-Jet*

Helmet streamers have been postulated to play a key role in the processes of jet formation in microquasars/AGN



( Gouveia dal Pino 2005 )

... and in outflows in young stellar objects.


( Ostriker & Shu 1995; Ferreira et al. 2006 ).

## Conclusions:

1. Helmet streamers are present also on fully / nearly fully convective objects.
2. Their discovery in stars other than the Sun (at a wavelength observable with high resolution) will therefore lead to a deeper understanding of these important magnetic structures.





Thank  
You!



First VLBI image of the system

Phillips et al. (1996)



At the short periastron distance the two coronae nearly overlap, giving rise to the observed giant flares.

The result of such a situation, where the whole streamer participates in the magnetic reconnection process, is a large structure, which holds the emission peaks at the position of the two helmets anchored on each of the two stars as in the VLBI image of Phillips and collaborators (1996).

Comparison of the Solar Corona at Solar Maximum and Minimum  
(White Light Eclipse Images from the High Altitude Observatory)



## SOLAR MAXIMUM

## SOLAR MINIMUM

Light seen during solar eclipses (Maximum and Minimum):

It does not come from coronal emission but is photospheric light scattered from the coronal gas